

Ferrofluid dynamics by ODD and EVEN Neutron Spin Echo studied

V.T.Lebedev¹, Gy.Torok², L.Vekas³

¹ Petersburg Nuclear Physics Institute, 188300 Gatchina, St.Petersburg distr., Russia

² Research Institute for Solid State Physics and Optics, Konkoly-Thege ut. 29-33, Budapest, Hungary

³ Center of Fundamental and Advanced Thechnical Research, Timisoara-1900, Romania

Odd neutron spin echo (ODD-NSE)[1] combined with NSE(EVEN) [2] yield the excitations spectra in condensed matter $S(\omega, \mathbf{q}) = S_{\text{odd}}(\omega, \mathbf{q}) + S_{\text{even}}(\omega, \mathbf{q})$ from odd and even NSE-signals dependent on time t and momentum transfer \mathbf{q} :

$$S_{\text{odd}} \sim \int S(\omega, \mathbf{q}) \sin(\omega t) d\omega,$$

$$S_{\text{even}} \sim \int S(\omega, \mathbf{q}) \cos(\omega t) d\omega.$$

In ferrofluids (FF) the frequencies $\omega \leq 1/\tau$ and relaxation times τ of soft modes characterise particles' interaction masked by viscous medium. We have studied the FF not magnetised (20°C, density $d = 1.69 \text{ g/cm}^3$; 20%vol. of Fe_3O_4 stabilised in pentanole by oleic and dodecylbenzenesulphone acids' bilayer; magnetisation $4\pi M_s = 860 \text{ G}$). Each particle (diameter $D_p \sim 16 \text{ nm}$, core $D_m \sim 10 \text{ nm}$), surrounded by other particles (spacing $\sim 20 \text{ nm} \sim D_p$), can move only by their regrouping (cage effect). Indeed, the NSE-data (Fig.1) confirm a delayed correlations $S_{\text{even}}(t)$ damping started at $t \gtrsim t^* \sim 10 \text{ ns}$ for momentum transfer $q_{1,2,3}$ (0.31 ; 0.55 ; 0.91 nm^{-1}). At $q > 1/D_p$ the even part shows a diffusion of single particle by cage slowed. The $S_{\text{odd}}(t) \sim t$ grow quite linear at $t \leq 1 \text{ ns}$, their slope gives the energy transfer q -dependent. At $t > 10 \text{ ns}$ the curves converge.

In gaussian approximation $S_{\text{even}}(t) \sim \exp[-q^2 \Gamma(t)/2]$ describes particle's self-correlation, where $\Gamma(t)$ is a squared particle's displacement for the time t . The $S_{\text{odd}}(t)$ do not adhere a simple model, $S_{\text{odd}}(t/\tau) \sim t \cdot \exp(-t/\tau)$. The normalised functions $F(t) = S_{\text{odd}}(t)/t$ display oscillations and damping at $q \sim 0.6 - 0.9 \text{ nm}^{-1}$ (Fig.2).




Figure 1: Even (1,3,5) and odd (2,4,6) functions: 20°C, $q_1(1,2)$; $q_2(3,4)$, $q_3(5,6)$. Lines are spline-functions.

Note, neutrons excite particles' vibrations:

$F(t) = \langle \omega \rangle + \beta \cdot t^2 + \mu \cdot \cos(\Omega t) \exp(-\nu t)$ where fitting parameters are q -dependent (Tab.1). The $q_1 \sim \pi/D_m$ is maximum wave number for acoustic wave, frequency $\Omega \sim 0.5 \text{ rad./ns}$, in coupled particles chain (spacing \sim core diameter $D_m \sim 10 \text{ nm}$). At $q_{2,3} > q_1$ the optical modes only permitted have frequency $\Omega \sim 0.3 \text{ rad./ns}$, period $T_\Omega \sim 20 \text{ ns}$, damping time $\tau = 1/\nu \sim 10 \text{ ns} \sim T_\Omega/2$.

The slow mode has frequencies $\langle \omega \rangle = \int \omega \cdot S(\omega, q) d\omega / \int S(\omega, q) d\omega \sim (3-5) \cdot 10^7 \text{ rad./s}$. The periods $T = 2\pi/\langle \omega \rangle \sim 100 \text{ ns}$ at $q_{2,3}$, and $T \sim 200 \text{ ns}$ at q_1 are much longer than the experimental time $t = 0 - 20 \text{ ns}$. The estimated velocity of slow waves propagation $U \sim \langle \omega \rangle / q \sim 10 \text{ cm/s}$ is by 4 orders in magnitude smaller than sound velocity in liquids.




Figure 2: Function $S_{\text{odd}}(t)/t$ at $q_{1,2,3}$ (1,2,3). Lines are fitting functions $F(t)$.

q , nm^{-1}	0.31	0.55	0.91
$\langle\omega\rangle$, ns^{-1}	0.029 ± 0.002	0.049 ± 0.002	0.051 ± 0.001
β , ns^{-3}	$(5\pm 1)\cdot 10^{-5}$	—	—
μ , ns^{-1}	-0.0044 ± 0.0011	0.017 ± 0.010	0.042 ± 0.011
Ω , ns^{-1}	0.50 ± 0.02	0.32 ± 0.02	0.30 ± 0.02
v , ns^{-1}	—	0.072 ± 0.062	0.14 ± 0.04

Tab. 1: Odd function: parameters.

The ratio $\mu/\langle\omega\rangle \sim 0.2-0.8$, growing with q , indicates the balance of fast and slow dynamics. The latter dominates at small q , when the odd-correlations are also delayed for $t^* \sim 10\text{ns}$, that is described by the term $\beta \cdot t^2$ in function $F(t)$. At the scale of radius $2\pi/q_3 \sim D_m/2$ fast and slow components in dynamics are comparable.

The $S(t)_{\text{even}} = \exp[-q^2\Gamma(t)/2]$ depend on particle's shift $\Delta X = \Gamma(t)^{1/2} = [-(2/q^2)\ln(S)]^{1/2}$ increasing slowly up to $\Delta X(t) \sim 1\text{nm}$ at $t < 15\text{ns}$. Then it achieves the $\Delta X \sim 3\text{nm} \sim$ radius of core (Fig.3). At $t < 15\text{ns}$ the shift adheres the function $\Delta X(t) = [a^2 + (vt)^2]^{1/2}$. Here $a = 0.4 \pm 0.1\text{nm}$ is the amplitude of fast oscillations, $v = 8.7 \pm 0.3\text{ cm/s}$ is the component of particle's velocity along q (data for $q = q_1$).




Figure 3: Particle's shift: data 1-3 at $q_{1,2,3}$, fitting function $\Delta X = [a^2 + (vt)^2]^{1/2}$ for data 1.

Vibrating with a small amplitude (\sim solvent molecule), a particle moves slowly waiting the conditions for a jump (length \sim shell thickness \sim core radius). Its velocity $v \sim 10\text{cm/s}$ is smaller by an order in magnitude than thermal one, $V_T = (kT/m_p)^{1/2} \sim 100\text{cm/s}$ (m_p is particle's mass). Hence, a particle involves solvent molecules and neighbouring particles (1st, 2nd spheres, region ~ 100 volumes of particle). Comparing even and odd dynamics, we conclude: the damping time of particle's oscillations $\tau = 1/v \sim T_Q/2 \sim 10\text{ns}$ is comparable with induction period, after that it jumps from cage to a new position.

Even data show particle's motion as a fast vibration + slow moving and final jump. Odd data visualise the main dynamic component that is a soft mode. Without it a jump of particle to new position is not possible. Odd NSE-signal reflects the potential defining particles motion. In FF with massive particles we observe their fast vibrations and relaxation initial stage, $S_{\text{odd}}(t) \sim t$.

References

- [1] Lebedev V.T., Torok Gy. Odd neutron spin echo // 3rd European Conf. on Neutron Scattering. Sept. 3-6, 2003, Montpellier, France. Abstract book, p.155.
- [2] Mezei F. Neutron Spin-Echo: a new concept in polarised neutron techniques. // Z.Physik, 1972, v.255, p.146-160.